

# *Ranunculus trilobus*

## Habitus and growth type

Plant height [m]: **0.22**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **1.02**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **7**

Reaction indicator value: **4.7**

Nutrient indicator value: **4.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.92**

Disturbance frequency (herb layer): **1.91**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.2**

Grazing pressure: **0.25**

Soil disturbance: **0.23**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*OC \(ISO\) \*Isoëto-Nanojuncetea\*, QC \(CHE\) \*Chenopodietea\*\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Synanthropic**