

Rhinanthus borbasisii

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **6**

Moisture indicator value: **3**

Reaction indicator value: **9**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.08**

Disturbance frequency (herb layer): **2.08**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **1.12**

Grazing pressure: **0.34**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**