

# ***Rhinanthus burnatii***

## **Habitus and growth type**

Plant height [m]: **0.26**

Life span: **Annual**

Life form: **Therophyte**

## **Flower**

Flowering period: **April-June**

## **Fruit, seed and dispersal**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## **Trophic mode**

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **7.5**

Temperature indicator value: **6**

Moisture indicator value: **5.3**

Reaction indicator value: **7**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.29**

Disturbance frequency (herb layer): **1.52**

Disturbance severity: **0.61**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0.12**

Grazing pressure: **0.27**

Soil disturbance: **0.13**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) Trifolio-Geranietea sanguinei](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**