

Rorippa brachycarpa

Habitus and growth type

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **4.8**

Moisture indicator value: **6**

Disturbance indicator values

Disturbance frequency: **1.7**

Disturbance frequency (herb layer): **2.01**

Disturbance severity: **0.56**

Disturbance severity (herb layer): **0.55**

Mowing frequency: **0.54**

Grazing pressure: **0.2**

Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**CM \(MOL\) *Molinio-Arrhenatheretea***](#),
[**DB \(FEP\) *Festuco-Puccinellietea***](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Saline vegetation**