

# *Rosa gallica*

## Habitus and growth type

Plant height [m]: **0.93**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **764.56**

Dispersal mode: **Endozoochory, Anthropochory**

Dispersal distance class: **6, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.2**

Temperature indicator value: **6.8**

Moisture indicator value: **4.1**

Reaction indicator value: **7.2**

Nutrient indicator value: **3.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.32**  
Disturbance frequency (herb layer): **1.47**  
Disturbance severity: **0.58**  
Disturbance severity (herb layer): **0.28**  
Mowing frequency: **0.22**  
Grazing pressure: **0.24**  
Soil disturbance: **0.12**

### **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) \*Trifolio-Geranietea sanguinei\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

### **Distribution**

Continentality: **6**  
Continentality amplitude: **3**