

# *Rosa sherardii*

## Habitus and growth type

Plant height [m]: **1.59**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Seed mass [mg]: **12.06**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **6.3**

Moisture indicator value: **5**

Reaction indicator value: **6.7**

Nutrient indicator value: **4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.81**  
Disturbance frequency (herb layer): **1.4**  
Disturbance severity: **0.5**  
Disturbance severity (herb layer): **0.33**  
Mowing frequency: **0.17**  
Grazing pressure: **0.23**  
Soil disturbance: **0.13**

### **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [CD \(RHA\) Crataego-Prunetea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

### **Distribution**

Continentality: **4**

Continentality amplitude: **4**