

# *Rosa spinosissima*

## Habitus and growth type

Plant height [m]: **0.77**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Specific leaf area [mm<sup>2</sup>/mg]: **15.62**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **20.41**

Dispersal mode: **Endozoochory, Anthropochory**

Dispersal distance class: **6, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **5.8**

Moisture indicator value: **4.1**

Reaction indicator value: **7.7**

Nutrient indicator value: **2.8**

Salinity indicator value: **0**

## Disturbance indicator values

Disturbance frequency: **0.57**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.17**

Grazing pressure: **0.26**

Soil disturbance: **0.16**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CD \(RHA\) \*Crataego-Prunetea\*](#), [CL \(GER\) \*Trifolio-Geranietea sanguinei\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [N1A Atlantic and Baltic coastal dune scrub](#)

Constant species of EUNIS habitats: [N1A Atlantic and Baltic coastal dune scrub](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),  
Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality amplitude: **8**