

Rubus caesius

Habitus and growth type

Plant height [m]: **0.68**

Life span: **Perennial**

Life form: **Phanerophyte, Chamaephyte, Shrub**

Leaf

Specific leaf area [mm²/mg]: **18.29**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **235.79**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Ellenberg-type indicator values

Light indicator value: **6.9**

Temperature indicator value: **5.3**

Moisture indicator value: **6.4**

Reaction indicator value: **7.3**

Nutrient indicator value: **6.7**

Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.37**

Disturbance frequency (herb layer): **1.4**

Disturbance severity: **0.6**

Disturbance severity (herb layer): **0.35**

Mowing frequency: **0.13**

Grazing pressure: **0.2**

Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**CF \(ROB\) Robinietea, HA \(POP\) Alno glutinosae-Populetea albae**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**N1A Atlantic and Baltic coastal dune scrub, N1D Atlantic and Baltic broad-leaved coastal dune forest, S32 Temperate Rubus scrub, T11 Temperate Salix and Populus riparian forest, T13 Temperate hardwood riparian forest**](#)

Constant species of EUNIS habitats: [**N1A Atlantic and Baltic coastal dune scrub, N1D Atlantic and Baltic broad-leaved coastal dune forest, S32 Temperate Rubus scrub, S35 Temperate and submediterranean thorn scrub, S91 Temperate riparian scrub, T11 Temperate Salix and Populus riparian forest, T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils, T13 Temperate hardwood riparian forest, T1H Broadleaved deciduous plantation of non site-native trees, V39 Mesic perennial anthropogenic herbaceous vegetation**](#)

Dominant species of EUNIS habitats: [**N1A Atlantic and Baltic coastal dune scrub, S32 Temperate Rubus scrub**](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **7**