

Ruta chalepensis

Habitus and growth type

Plant height [m]: **0.6**

Life span: **Perennial**

Life form: **Chamaephyte**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Seed mass [mg]: **1.32**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.6**

Temperature indicator value: **9**

Moisture indicator value: **2.3**

Reaction indicator value: **7.6**

Nutrient indicator value: **4.8**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.47**

Disturbance frequency (herb layer): **1.42**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.07**

Grazing pressure: **0.34**

Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) *Ononido-Rosmarinetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S54 *Thermomediterranean arid scrub*](#)

Constant species of EUNIS habitats: [S54 *Thermomediterranean arid scrub*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**