

Sagina nodosa

Habitus and growth type

Plant height [m]: **0.12**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **15.5**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.03**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7.9**

Temperature indicator value: **5**

Moisture indicator value: **7.7**

Reaction indicator value: **7**

Nutrient indicator value: **4.3**

Salinity indicator value: **0.8**

Disturbance indicator values

Disturbance frequency: **1.79**

Disturbance frequency (herb layer): **1.96**
Disturbance severity: **0.47**
Disturbance severity (herb layer): **0.47**
Mowing frequency: **0.28**
Grazing pressure: **0.44**
Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [JA \(SAG\) *Saginetea maritima*](#), [OC \(ISO\) *Isoëto-Nanojuncetea*](#), [PA \(SCH\) *Scheuchzerio palustris-Caricetea fuscae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [MA222 Atlantic upper saltmarsh](#)

Constant species of EUNIS habitats: [MA222 Atlantic upper saltmarsh](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Spring, Wetland, Mire, Saline vegetation**

Distribution

Continentality amplitude: **6**