

# *Salix myrsinifolia*

## Habitus and growth type

Plant height [m]: **3.33**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Specific leaf area [mm<sup>2</sup>/mg]: **11.27**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.09**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **3.6**

Moisture indicator value: **7.6**

Reaction indicator value: **6.3**

Nutrient indicator value: **4.5**

Salinity indicator value: **0**

## Disturbance indicator values

Disturbance frequency: **0.51**

Disturbance frequency (herb layer): **1.48**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.11**

Grazing pressure: **0.18**

Soil disturbance: **0.13**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*U25 Boreal and Arctic base-rich scree and block field\*\*](#)

Constant species of EUNIS habitats: [\*\*U25 Boreal and Arctic base-rich scree and block field\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Alpine-subalpine and arctic grassland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **5**

Continentality amplitude: **4**