

# *Salix pentandra*

## Habitus and growth type

Plant height [m]: **8.53**

Life span: **Perennial**

Life form: **Phanerophyte, Tree, Shrub**

## Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Specific leaf area [mm<sup>2</sup>/mg]: **18.42**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.13**

Dispersal mode: **Anemochory, Anthropochory**

Dispersal distance class: **5, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **4.5**

Moisture indicator value: **8.1**

Reaction indicator value: **5.7**

Nutrient indicator value: **4**

Salinity indicator value: **0**

## Disturbance indicator values

Disturbance frequency: **0.68**

Disturbance frequency (herb layer): **1.33**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.22**

Mowing frequency: **0.13**

Grazing pressure: **0.17**

Soil disturbance: **0.18**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CR \(VIR\) \*Betulo carpaticae-Alnetea viridis\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S92 \*Salix fen scrub\*](#)

### Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **6**

Continentality amplitude: **5**