

# ***Salix phylicifolia***

## **Habitus and growth type**

Plant height [m]: **1.5**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **10.53**

## **Fruit, seed and dispersal**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **3.3**

Moisture indicator value: **8.1**

Reaction indicator value: **5**

Nutrient indicator value: **4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.59**

Disturbance frequency (herb layer): **1.38**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.22**

Mowing frequency: **0.15**

Grazing pressure: **0.15**

Soil disturbance: **0.15**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **[CR \(VIR\) Betulo carpaticae-Alnetea viridis](#)**

EUNIS habitat

Diagnostic species of EUNIS habitats: **[U35 Boreal and Arctic base-rich inland cliff](#)**

Constant species of EUNIS habitats: **[U35 Boreal and Arctic base-rich inland cliff](#)**

Broad habitat

Occurrence in broad habitats: **Scrub, Sparsely vegetated (incl. rock and scree)**