Salix salviifolia

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **6**
Moisture indicator value: **12**
Reaction indicator value: **6**
Nutrient indicator value: **5**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.62**
Disturbance frequency (herb layer): **1.66**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.38**
Mowing frequency: **0.01**
Grazing pressure: **0.24**
Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FD (NER) *Nerio-Tamaricetea*, HB (PUR) *Salicetea purpureae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **U72 Unvegetated or sparsely vegetated gravel**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

bar in the Mediterranean region

Broad habitat

Occurrence in broad habitats: **Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz