

Salix viminalis

Habitus and growth type

Plant height [m]: **6.08**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Specific leaf area [mm²/mg]: **12.06**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **5.8**

Moisture indicator value: **8**

Reaction indicator value: **7.1**

Nutrient indicator value: **5.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.78**
Disturbance frequency (herb layer): **1.59**
Disturbance severity: **0.59**
Disturbance severity (herb layer): **0.43**
Mowing frequency: **0.08**
Grazing pressure: **0.23**
Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [HB \(PUR\) *Salicetea purpureae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S91 Temperate riparian scrub](#), [T11 Temperate *Salix* and *Populus* riparian forest](#)

Constant species of EUNIS habitats: [S91 Temperate riparian scrub](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **6**