

# *Salvia aethiopis*

## Habitus and growth type

Plant height [m]: **0.74**

Life span: **Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **1.75**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.4**

Temperature indicator value: **7.1**

Moisture indicator value: **3.4**

Reaction indicator value: **7.5**

Nutrient indicator value: **4.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.51**

Disturbance frequency (herb layer): **1.88**

Disturbance severity: **0.61**

Disturbance severity (herb layer): **0.58**

Mowing frequency: **0.76**

Grazing pressure: **0.33**

Soil disturbance: **0.26**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [QF \(ART\) Artemisietea vulgaris](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**