

# *Salvia verticillata*

## Habitus and growth type

Plant height [m]: **0.47**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **15.1**

## Flower

Flowering period: **May-September**

## Fruit, seed and dispersal

Seed mass [mg]: **0.7**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.9**

Temperature indicator value: **5.9**

Moisture indicator value: **4.2**

Reaction indicator value: **7**

Nutrient indicator value: **5.2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.45**

Disturbance frequency (herb layer): **1.76**  
Disturbance severity: **0.52**  
Disturbance severity (herb layer): **0.5**  
Mowing frequency: **0.77**  
Grazing pressure: **0.26**  
Soil disturbance: **0.18**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [QF \(ART\) Artemisietea vulgaris](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#)

Constant species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

## Distribution

Continentality: **5**  
Continentality amplitude: **4**