Sanguisorba verrucosa

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **8.48**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **8**
Moisture indicator value: **3**
Nutrient indicator value: **2.7**
Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **0.56**
Disturbance frequency (herb layer): **1.66**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.28**
Grazing pressure: **0.33**
Soil disturbance: **0.25**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FG (SAC) *Stipo giganteae-Agrostietea castellanae*, FJ (TRA) *Stipo-Trachynietea distachyae***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz