Saxifraga federici-augusti

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.9**
Temperature indicator value: **7.3**
Moisture indicator value: **2.3**
Reaction indicator value: **7.4**
Nutrient indicator value: **2.4**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.65**
Disturbance frequency (herb layer): **1.74**
Disturbance severity: **0.39**
Disturbance severity (herb layer): **0.26**
Mowing frequency: **0**
Grazing pressure: **0.27**
Soil disturbance: **0.13**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz