

# *Saxifraga pedemontana*

## Habitus and growth type

Plant height [m]: **0.07**

Life span: **Perennial**

Life form: **Chamaephyte**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.04**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **3.5**

Temperature indicator value: **2.5**

Moisture indicator value: **3**

Reaction indicator value: **3.5**

Nutrient indicator value: **1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.86**

Disturbance frequency (herb layer): **1.01**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.01**

Grazing pressure: **0.13**

Soil disturbance: **0.11**

## **Habitat and sociology**

EUNIS habitat

Diagnostic species of EUNIS habitats: [U24 Mediterranean siliceous scree](#), [U34 Mediterranean siliceous inland cliff](#)

Constant species of EUNIS habitats: [U24 Mediterranean siliceous scree](#), [U34 Mediterranean siliceous inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**