

# *Saxifraga umbrosa*

## Habitus and growth type

Plant height [m]: **0.26**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **3**

Temperature indicator value: **5**

Moisture indicator value: **5**

Reaction indicator value: **3**

Nutrient indicator value: **4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.13**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.16**

Mowing frequency: **0.03**

Grazing pressure: **0.2**

Soil disturbance: **0.09**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [BA \(PIC\) Vaccinio-Piceetea](#), [CA \(FAG\) Carpino-Fagetea sylvaticae](#)

Broad habitat

Occurrence in broad habitats: **Forest**