Scaligeria napiformis

Habitus and growth type

Life span: **Biennial or short-lived**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **1.36**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Moisture indicator value: **4**
Nutrient indicator value: **3**
Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.21**
Disturbance frequency (herb layer): **1.5**
Disturbance severity: **0.72**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.04**
Grazing pressure: **0.32**
Soil disturbance: **0.17**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S63 Eastern garrigue**, **T3D Mediterranean Cupressaceae forest**

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz