

# *Scorpiurus vermiculatus*

## Habitus and growth type

Plant height [m]: **0.13**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **March-April**

## Fruit, seed and dispersal

Seed mass [mg]: **6.52**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **8**

Moisture indicator value: **2.5**

Reaction indicator value: **6**

Nutrient indicator value: **4**

Salinity indicator value: **0**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FH \(BUL\) \*Poetea bulbosae\*](#), [QC \(CHE\) \*Chenopodietea\*](#)

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**