

# *Scorzonera purpurea*

## Habitus and growth type

Plant height [m]: **0.33**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.54**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.8**

Temperature indicator value: **6.6**

Moisture indicator value: **3.1**

Reaction indicator value: **8**

Nutrient indicator value: **2.2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.26**

Disturbance frequency (herb layer): **1.93**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.77**

Grazing pressure: **0.34**

Soil disturbance: **0.14**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [FP \(ONO\) Festuco hystricis-Ononidetea striatae](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## **Distribution**

Continentality: **8**

Continentality amplitude: **5**