Scorzoneroides cantabrica

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **3**
Moisture indicator value: **3**
Reaction indicator value: **2**
Nutrient indicator value: **1**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.34**
Disturbance frequency (herb layer): **1.55**
Disturbance severity: **0.35**
Disturbance severity (herb layer): **0.32**
Mowing frequency: **0.08**
Grazing pressure: **0.27**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FL (IND) *Festucetea indigestae***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz