Scrophularia nodosa

Habitus and growth type

Plant height [m]: **0.69**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]:

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.11**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.8**
Temperature indicator value: **5.1**
Moisture indicator value: **6.3**
Reaction indicator value: **6.6**
Nutrient indicator value: **6.9**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.13**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.35**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.24**
Mowing frequency: **0.07**
Grazing pressure: **0.19**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae*, HA (POP) *Alno glutinosae-Populetea albae***

EUNIS habitat

Constant species of EUNIS habitats: **T13 Temperate hardwood riparian forest, T1E Carpinus and Quercus mesic deciduous forest**

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz