Scrophularia vernalis

Habitus and growth type

Plant height [m]: **0.6**
Life span: **Biennial or short-lived, Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **23.8**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.19**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.9**
Temperature indicator value: **5.8**
Moisture indicator value: **5.4**
Reaction indicator value: **6.7**
Nutrient indicator value: **5.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.1**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.27**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.19**
Mowing frequency: **0.03**
Grazing pressure: **0.16**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae***

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz