

Scutellaria minor

Habitus and growth type

Plant height [m]: **0.14**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **37.31**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.2**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**

Temperature indicator value: **6**

Moisture indicator value: **9**

Reaction indicator value: **2.6**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.03**

Disturbance frequency (herb layer): **1.55**
Disturbance severity: **0.38**
Disturbance severity (herb layer): **0.32**
Mowing frequency: **0.34**
Grazing pressure: **0.21**
Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#),
[PA \(SCH\) *Scheuchzerio palustris-Caricetea fuscae*](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Mire, Grassland (non-alpine, non-saline)**

Distribution

Continentality: **2**
Continentality amplitude: **2**