

Sedum hispanicum

Habitus and growth type

Plant height [m]: **0.08**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.03**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.6**

Moisture indicator value: **2.7**

Reaction indicator value: **5.3**

Nutrient indicator value: **2.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.74**

Disturbance frequency (herb layer): **1.61**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.44**

Mowing frequency: **0.27**

Grazing pressure: **0.3**

Soil disturbance: **0.24**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) *Sedo-Scleranthetea*, KC \(ASP\) *Asplenietea trichomanis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Constant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**