

Schoenoplectus triqueter

Habitus and growth type

Plant height [m]: **0.74**

Life span: **Perennial**

Life form: **Geophyte**

Leaf

Specific leaf area [mm²/mg]: **6.84**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **1.15**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **7**

Moisture indicator value: **9.7**

Reaction indicator value: **7**

Nutrient indicator value: **7**

Salinity indicator value: **1.8**

Disturbance indicator values

Disturbance frequency: **1.81**

Disturbance frequency (herb layer): **1.81**
Disturbance severity: **0.29**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.13**
Grazing pressure: **0.07**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OD \(PHR\) *Phragmito-Magnocaricetea*](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Aquatic, Wetland, Saline vegetation**

Distribution

Continentality amplitude: **9**