

Silene viscaria

Habitus and growth type

Plant height [m]: **0.42**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **14.29**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.07**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Moisture indicator value: **3.5**

Reaction indicator value: **4**

Nutrient indicator value: **2.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.4**

Disturbance frequency (herb layer): **1.58**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.33**

Mowing frequency: **0.32**

Grazing pressure: **0.22**

Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) *Trifolio-Geranietea sanguinei*](#), [DA \(FES\) *Festuco-Brometea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3H *Larix light taiga*](#)

Constant species of EUNIS habitats: [T3H *Larix light taiga*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **5**

Continentality amplitude: **4**