

# *Spergularia macrorrhiza*

## Habitus and growth type

Plant height [m]: **0.09**

Life span: **Perennial**

Life form: **Chamaephyte**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.04**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **10**

Moisture indicator value: **2**

Reaction indicator value: **6**

Nutrient indicator value: **1**

Salinity indicator value: **3**

### Disturbance indicator values

Disturbance frequency: **0.94**

Disturbance frequency (herb layer): **0.94**

Disturbance severity: **0.28**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.05**

Grazing pressure: **0.11**

Soil disturbance: **0.12**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [JA \(SAG\) \*Saginetea maritima\*](#), [JB \(CRI\) \*Crithmo-Staticetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [MA251 Mediterranean upper saltmarsh](#)

### Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal beach, dune or shingle, Coastal cliff, Saline vegetation, Sparsely vegetated (incl. rock and scree)**