

Spergularia rubra

Habitus and growth type

Plant height [m]: **0.13**

Life span: **Annual, Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm²/mg]: **14.64**

Flower

Flowering period: **February-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.04**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**

Moisture indicator value: **4.6**

Reaction indicator value: **4**

Nutrient indicator value: **3.9**

Salinity indicator value: **0.4**

Disturbance indicator values

Disturbance frequency: **1.79**

Disturbance frequency (herb layer): **2.1**

Disturbance severity: **0.73**

Disturbance severity (herb layer): **0.72**

Mowing frequency: **0.4**

Grazing pressure: **0.22**

Soil disturbance: **0.54**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Aquatic, Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: 7