

Stachys alopecuroides

Habitus and growth type

Plant height [m]: **0.26**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **22.2**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.19**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **3**

Moisture indicator value: **4.4**

Reaction indicator value: **7.5**

Nutrient indicator value: **2.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.28**

Disturbance frequency (herb layer): **0.92**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.22**
Mowing frequency: **0.08**
Grazing pressure: **0.22**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) *Elyno-Seslerietea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Constant species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**
Continentality amplitude: **1**