

Stachys arvensis

Habitus and growth type

Plant height [m]: **0.19**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **29.8**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.7**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**

Moisture indicator value: **4.6**

Nutrient indicator value: **5.9**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.31**

Disturbance frequency (herb layer): **1.9**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.65**

Mowing frequency: **0.68**

Grazing pressure: **0.24**

Soil disturbance: **0.54**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) *Papaveretea rhoeadis*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **3**

Continentality amplitude: **3**