

# *Stellaria palustris*

## Habitus and growth type

Plant height [m]: **0.26**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **23.02**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.48**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6**

Temperature indicator value: **5.2**

Moisture indicator value: **8.3**

Reaction indicator value: **5**

Nutrient indicator value: **3.2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.37**

Disturbance frequency (herb layer): **1.74**

Disturbance severity: **0.4**

Disturbance severity (herb layer): **0.38**

Mowing frequency: **0.55**

Grazing pressure: **0.21**

Soil disturbance: **0.14**

## **Habitat and sociology**

### **Syntaxon**

Diagnostic species of phytosociological classes: [OD \(PHR\) \*Phragmito-Magnocaricetea\*](#)

### **Broad habitat**

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire, Grassland (non-alpine, non-saline)**

## **Distribution**

Continentality: **6**

Continentality amplitude: **6**