

# *Styrax officinalis*

## Habitus and growth type

Plant height [m]: **4.75**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **411.55**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **9**

Reaction indicator value: **6.7**

Nutrient indicator value: **3.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.08**

Disturbance frequency (herb layer): **1.37**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.17**

Mowing frequency: **0**

Grazing pressure: **0.25**

Soil disturbance: **0.11**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#)

### Broad habitat

Occurrence in broad habitats: **Scrub, Forest**