Symphoricarpos albus

Habitus and growth type

Plant height [m]: **1.27**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**
Leaf phenology (woody plants only): **Winter deciduous**
Specific leaf area [mm^2/mg]: **19.36**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **6.99**
Dispersal mode: **Endozoochory, Anthropochory**
Dispersal distance class: **6, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6**
Temperature indicator value: **5.8**
Moisture indicator value: **5.3**
Reaction indicator value: **6**
Nutrient indicator value: **6.3**
Salinity indicator value: **0**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance indicator values

Disturbance frequency: **0.14**
Disturbance frequency (herb layer): **1.36**
Disturbance severity: **0.71**
Disturbance severity (herb layer): **0.22**
Mowing frequency: **0.05**
Grazing pressure: **0.19**
Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **HA (POP) *Alno glutinosae-Populetea albae***

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Synanthropic**

Distribution

Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz