

Symphyotrichum squamatum

Habitus and growth type

Plant height [m]: **0.81**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **August-October**

Fruit, seed and dispersal

Dispersal mode: **Anemochory, Anthropochory**

Dispersal distance class: **5, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **9**

Moisture indicator value: **7**

Reaction indicator value: **7.5**

Nutrient indicator value: **7.5**

Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **1.25**

Disturbance frequency (herb layer): **1.67**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.62**

Mowing frequency: **0.44**

Grazing pressure: **0.22**

Soil disturbance: **0.33**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**FF \(LYG\) *Lygeo sparti-Stipetea tenacissimae***](#), [**QF \(ART\) *Artemisietea vulgaris***](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**