Tamarix ramosissima

Habitus and growth type

Plant height [m]:
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Scale-like**
Leaf phenology (woody plants only): **Winter deciduous**

Fruit, seed and dispersal

Seed mass [mg]: **0.15**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**

Ellenberg-type indicator values

Light indicator value: **8.3**
Moisture indicator value: **7.2**
Reaction indicator value: **7.5**
Nutrient indicator value: **6**
Salinity indicator value: **3.5**

Disturbance indicator values

Disturbance frequency: **1.41**
Disturbance frequency (herb layer): **1.81**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.45**
Mowing frequency: **0.13**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.34**
Soil disturbance: **0.25**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CD (RHA) *Crataego-Prunetea*, EB (TAM) *Tamaricetea arceuthoidis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S94 Semi-desert riparian scrub**
Constant species of EUNIS habitats: **S94 Semi-desert riparian scrub**
Dominant species of EUNIS habitats: **S94 Semi-desert riparian scrub**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz