

Taraxacum besarabicum

Habitus and growth type

Plant height [m]: **0.13**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **6.4**

Moisture indicator value: **5.9**

Reaction indicator value: **8**

Nutrient indicator value: **4.5**

Salinity indicator value: **5.7**

Disturbance indicator values

Disturbance frequency: **2.18**

Disturbance frequency (herb layer): **2.18**

Disturbance severity: **0.45**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.6**

Grazing pressure: **0.55**

Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**DB \(FEP\) Festuco-Puccinellietea**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R63 Temperate inland salt marsh**](#)

Constant species of EUNIS habitats: [**R63 Temperate inland salt marsh**](#)

Broad habitat

Occurrence in broad habitats: **Saline vegetation**