Taraxacum besarabicum

Habitus and growth type

Plant height [m]: **0.13**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**

Ellenberg-type indicator values

Light indicator value: **8.7**
Temperature indicator value: **6.4**
Moisture indicator value: **5.9**
Reaction indicator value: **8**
Nutrient indicator value: **4.5**
Salinity indicator value: **5.7**

Disturbance indicator values

Disturbance frequency: **2.18**
Disturbance frequency (herb layer): **2.18**
Disturbance severity: **0.45**
Disturbance severity (herb layer): **0.45**
Mowing frequency: **0.6**
Grazing pressure: **0.55**
Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic species of phytosociological classes: **DB (FEP) *Festuco-Puccinellietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R63 Temperate inland salt marsh**

Constant species of EUNIS habitats: **R63 Temperate inland salt marsh**

Broad habitat

Occurrence in broad habitats: **Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz