

# ***Taraxacum besarabicum***

## **Habitus and growth type**

Plant height [m]: **0.13**

Life form: **Hemicryptophyte**

## **Fruit, seed and dispersal**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

### Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **6.4**

Moisture indicator value: **5.9**

Reaction indicator value: **8**

Nutrient indicator value: **4.5**

Salinity indicator value: **5.7**

### Disturbance indicator values

Disturbance frequency: **2.18**

Disturbance frequency (herb layer): **2.18**

Disturbance severity: **0.45**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.6**

Grazing pressure: **0.55**

Soil disturbance: **0.22**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*DB \(FEP\) Festuco-Puccinellietea\*\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*R63 Temperate inland salt marsh\*\*](#)

Constant species of EUNIS habitats: [\*\*R63 Temperate inland salt marsh\*\*](#)

Broad habitat

Occurrence in broad habitats: **Saline vegetation**