Taraxacum stevenii

Habitus and growth type

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **28.8**

Fruit, seed and dispersal

Seed mass [mg]: **1**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Disturbance indicator values

Disturbance frequency: **1.71**
Disturbance frequency (herb layer): **1.71**
Disturbance severity: **0.25**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.01**
Grazing pressure: **0.18**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **KF (LAM) *Lamio tomentosi-Chaerophylletea humilis*, LA (HER) *Salicetea herbaceae***

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz