

# *Teucrium expassum*

## Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

Spinescence: **not spinescent**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **6**

Moisture indicator value: **1**

Reaction indicator value: **8**

Nutrient indicator value: **1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.12**

Disturbance frequency (herb layer): **1.79**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.39**

Mowing frequency: **0.37**

Grazing pressure: **0.43**

Soil disturbance: **0.23**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) \*Ononido-Rosmarinetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1H Iberian oromediterranean basiphilous](#)

**dry grassland**

Constant species of EUNIS habitats: **R1H Iberian oromediterranean basiphilous dry grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**