Teucrium marum

Habitus and growth type

Plant height [m]: **0.4**
Life span: **Perennial**
Life form: **Chamaephyte**

Leaf

Leaf type (woody plants only): **Sclerophyllous**
Leaf phenology (woody plants only): **Evergreen**

Flower

Flowering period: **April-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.58**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **7.3**
Moisture indicator value: **3**
Reaction indicator value: **4**
Nutrient indicator value: **2.5**
Salinity indicator value: **0**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.72**
Disturbance frequency (herb layer): **1.33**
Disturbance severity: **0.63**
Disturbance severity (herb layer): **0.39**
Mowing frequency: **0.12**
Grazing pressure: **0.3**
Soil disturbance: **0.14**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S71 Western Mediterranean spiny heath**
Constant species of EUNIS habitats: **S71 Western Mediterranean spiny heath**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz