Thalictrum alpinum

Habitus and growth type

Plant height [m]: **0.08**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **14.35**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.75**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **2.5**
Moisture indicator value: **6.6**
Reaction indicator value: **5.8**
Nutrient indicator value: **2.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.47**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.65**
Disturbance severity: **0.27**
Disturbance severity (herb layer): **0.24**
Mowing frequency: **0.13**
Grazing pressure: **0.17**
Soil disturbance: **0.07**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CU (SES) *Elyno-Seslerietea*, PA (SCH) *Scheuchzerio palustris-Caricetea fuscae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q45 Arctic-alpine rich fen, R42 Boreal and Arctic acidophilous alpine grassland, U21 Boreal and Arctic siliceous scree and block field, U25 Boreal and Arctic base-rich scree and block field, U35 Boreal and Arctic base-rich inland cliff**

Constant species of EUNIS habitats: **Q45 Arctic-alpine rich fen, R42 Boreal and Arctic acidophilous alpine grassland, U21 Boreal and Arctic siliceous scree and block field, U25 Boreal and Arctic base-rich scree and block field, U35 Boreal and Arctic base-rich inland cliff**

Broad habitat

Occurrence in broad habitats: **Mire, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz