

Thalictrum minus

Habitus and growth type

Plant height [m]: **0.68**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **17.02**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **1.96**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.7**

Moisture indicator value: **3.2**

Reaction indicator value: **7.7**

Nutrient indicator value: **2.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.73**

Disturbance frequency (herb layer): **1.67**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.4**

Mowing frequency: **0.41**

Grazing pressure: **0.29**

Soil disturbance: **0.18**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#), [S36 Low steppic scrub](#), [T3H Larix light taiga](#)

Constant species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#), [S36 Low steppic scrub](#), [T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest](#), [T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest](#), [T3H Larix light taiga](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **8**