

# *Thesium linophyllon*

## Habitus and growth type

Plant height [m]: **0.34**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **15.52**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **4.77**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **6.1**

Moisture indicator value: **3.1**

Reaction indicator value: **8**

Nutrient indicator value: **1.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.78**

Disturbance frequency (herb layer): **1.58**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.4**

Mowing frequency: **0.47**

Grazing pressure: **0.28**

Soil disturbance: **0.13**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**