

Torilis leptophylla

Habitus and growth type

Plant height [m]: **0.25**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **23.64**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **1.37**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**

Temperature indicator value: **9.3**

Moisture indicator value: **3**

Reaction indicator value: **7.5**

Nutrient indicator value: **5.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.77**

Disturbance frequency (herb layer): **1.91**

Disturbance severity: **0.8**

Disturbance severity (herb layer): **0.66**

Mowing frequency: **0.39**

Grazing pressure: **0.25**

Soil disturbance: **0.41**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) *Stipo-Trachynietea distachyae*](#), [QC \(CHE\) *Chenopodietea*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**